


Size and structure of near-threshold states



Tetsuo Hyodo



TOKYO METROPOLITAN UNIVERSITY


東京都立大学

学問の力で、東京から世界の未来を拓く

2020, May. 18th₁

Hadron clusters

Hadrons near an s-wave two-body threshold



“hadronic molecules” (various flavors, baryon numbers, ...)

Two-body universal physics

Near-threshold s-wave state: universal physics

E. Braaten, H.-W. Hammer, Phys. Rept. 428, 259 (2006);

P. Naidon, S. Endo, Rept. Prog. Phys. 80, 056001 (2017)

- scattering length $|a| \gg$ interaction range R_0

- size of (quasi-)bound state $\sim |a|$: loosely bound

- relation with eigenenergy E


$$a(E) \sim \frac{i}{k} = \frac{i}{\sqrt{2\mu E}}$$

vdW

Examples: d , $\Lambda(1405)$, ${}^4\text{He}$ dimer

	NN [fm]	$\bar{K}N$ [fm]	${}^4\text{He}$ [a_0]
$a(E)$	4.3	1.2-0.8i	178
a_{emp}	5.1	1.4-0.9i	189
R_0	1.4	0.4	10

strong



${}^4\text{He}$

Wave function

Why is a weakly bound s-wave state spatially large?

- Radial Schrödinger equation

$$-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} u_0(r) + V(r)u_0(r) = Eu_0(r), \quad \psi_0(r) = Y_0^0 \frac{u_0(r)}{r}$$

- at large distance ($V(r) = 0$) with zero energy ($E = 0$)

$$u_0(r) = C(r - a), \quad (r > R_0)$$


- scattering length a : intercept of $u_0(r)$

- bound state with $B = 0 \Rightarrow |a| = \infty$

Wave function is not normalizable

$$\int d^3r |\psi_0(r)|^2 = \int_0^\infty dr |u_0(r)|^2 = \infty$$

$B = 0$ state is not a bound state (zero-energy resonance)



Consequences

Mean squared radius

$$\langle r^2 \rangle = \int d^3r \ r^2 |\psi_0(r)|^2 = \int_0^\infty dr \ r^2 |u_0(r)|^2 = \infty$$

—> size of $B = 0$ state is **infinitely large**

Compositeness X (weight of the composite component)

T. Hyodo, Phys. Rev. C 90, 055208 (2014)

$$\Psi = \begin{pmatrix} \tilde{\psi}_0(p) \\ c \\ \vdots \end{pmatrix} \quad X = \int \frac{d^3p}{(2\pi)^3} |\tilde{\psi}_0(p)|^2 = \int d^3r |\psi_0(r)|^2 \quad \text{infinite}$$

$$Z = |c|^2 + \dots \quad \text{finite}$$

$$X + Z = 1$$

—> $B = 0$ state is **completely composite** ($X = 1$, $Z = 0$)

Weakly bound state ($B \neq 0$, except for fine tuning)

- large spatial size and composite dominance

p-wave state

What about p-wave states?

- Radial Schrödinger equation

$$-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} u_1(r) + V(r)u_1(r) + \frac{\hbar^2}{\mu r^2} u_1(r) = Eu_1(r), \quad \psi_1(\mathbf{r}) = Y_1^m(\Omega) \frac{u_1(r)}{r}$$

- at large distance ($V(r) = 0$) with zero energy ($E = 0$)

$$u_1(r) = Cr^s, \quad s = 2, -1$$


- bound state with $B = 0$

$$u_1(r) = \frac{C}{r}, \quad (r > R_0)$$

Wave function is normalizable

$$\int d^3r |\psi_1(\mathbf{r})|^2 = \int_0^\infty dr |u_1(r)|^2 < \infty$$

$B = 0$ state is a bound state



Consequences (p-wave)

Mean squared radius

$$\langle r^2 \rangle = \int_0^\infty dr \ r^2 |u_1(r)|^2 = A_{\text{in}} + |C|^2 \int_{R_0}^\infty dr \ 1 = \infty \quad \leftarrow u_1(r) = \frac{C}{r}$$

—> size of $B = 0$ state is infinitely large

Compositeness X

T. Hyodo, Phys. Rev. C 90, 055208 (2014)

$$\Psi = \begin{pmatrix} \tilde{\psi}_1(\mathbf{p}) \\ c \\ \vdots \end{pmatrix} \quad X = \int \frac{d^3 p}{(2\pi)^3} |\tilde{\psi}_1(\mathbf{p})|^2 = \int d^3 r |\psi_1(\mathbf{r})|^2 \quad \text{finite}$$

$$Z = |c|^2 + \dots \quad \text{finite}$$

$$X + Z = 1$$

—> $B = 0$ can have any structure ($0 < X < 1$)

Weakly bound p-wave state ($B \neq 0$, except for fine tuning)

- large spatial size?

Higher partial waves

Wave function of a $B = 0$ state with angular momentum ℓ

$$u_\ell(r) = \frac{C}{r^\ell} \quad (r > R_0)$$

- **Mean squared radius**

$$\langle r^2 \rangle = \int_0^\infty dr \ r^2 |u_\ell(r)|^2 = A_{\text{in}} + |C|^2 \int_{R_0}^\infty dr \ r^{2-2\ell} = \infty \quad \text{if } \ell \leq 1$$

—> diverges only for s- and p-waves

Generalized radius


$$\sqrt[n]{\langle r^n \rangle} = \sqrt[n]{\int d^3r \ r^n |\psi(r)|^2} = \infty \quad \text{if } n - 2\ell \geq 0$$

- **s-wave: radius diverges for all $n \geq 0$** ← universality
- **p-wave: radius diverges for $n \geq 2$**
- **ℓ -th wave: radius diverges for $n \geq 2\ell$**

example: square well potential

Square well potential

$$V(r) = \begin{cases} -V_0 & 0 \leq r \leq b \\ 0 & b < r \end{cases}$$



- s-wave

$$\langle r^2 \rangle = \left[\frac{b}{2} - \frac{\sin(2kb)}{4k} + \frac{\sin^2(kb)}{2\kappa} \right]^{-1} \left[\frac{1}{k^3} \left(\frac{(kb)^3}{6} + \frac{1 - 2(kb)^2}{8} \sin(2kb) - \frac{kb}{4} \cos(2kb) \right) + \frac{\sin^2(kb)}{2\kappa^3} \left[(\kappa b)^2 + \kappa b + \frac{1}{2} \right] \right]$$

$$k = \frac{\sqrt{2\mu(V_0 - B)}}{\hbar}, \quad \kappa = \frac{\sqrt{2\mu B}}{\hbar}$$



- p-wave

$$\langle r^2 \rangle = \left[\frac{b}{2} + \frac{\sin(2kb)}{4k} - \frac{\sin^2(kb)}{k^2 b} + \frac{\kappa (\sin(kb) - kb \cos(kb))^2}{2k^2 (1 + \kappa b)^2} \left(1 + \frac{2}{\kappa b} \right) \right]^{-1}$$

$$\times \left[\frac{1}{k^3} \left(\frac{3kb + (kb)^3}{6} + \frac{-5 + 2(kb)^2}{8} \sin(2kb) + \frac{3kb}{4} \cos(2kb) \right) + \frac{(\sin(kb) - kb \cos(kb))^2}{2\kappa k^2 (1 + \kappa b)^2} \left[(\kappa b)^2 + 3\kappa b + \frac{5}{2} \right] \right]$$

Demonstration

Root mean squared radius $\sqrt{\langle r^2 \rangle}$ v.s. $\kappa = \sqrt{2\mu B}/\hbar$



Both radii diverge in the weak binding limit ($\kappa \rightarrow 0$).

Dependence on b (well width) seems different.

Weak-binding behavior

Behavior near the weak-binding limit

- s-wave

$$\langle r^2 \rangle = \frac{1}{2\kappa^2} + \mathcal{O}(\kappa^{-1})$$



independent of b (well width)

← universality

- p-wave

$$\langle r^2 \rangle = \frac{5b}{6} \frac{1}{\kappa} + \mathcal{O}(\kappa^0)$$

depends on b (well width)
non-universal divergence



Summary



Size and structure of $B = 0$ states

	$\langle r^2 \rangle$	compositeness	$\int d^3r \psi ^2$
s-wave	∞ (universal)	$X = 1$	∞
p-wave	∞ (non-universal)	$0 < X < 1$	finite



Implication: large mean squared radius of near-threshold ($B \neq 0$) p-wave bound states



What about resonances?

T. Hyodo, in preparation